

State Inventory Tool

User Guide for Estimating Emissions and Sinks from Land Use, Land-Use Change, and Forestry

SEPTEMBER 2026

ACKNOWLEDGEMENTS

The SIT was originally developed in conjunction with EPA’s state and local programs to automate the steps states would need to take in developing their own emission estimates in a manner that was consistent with prevailing national and state guidelines. The result was a user-friendly set of eleven modules that help users estimate greenhouse gas emissions and removals at the state level. CGS is supporting updates to the tool to facilitate continuity in state inventory development and improvement work at the state level.

Modules and User Guides updates prepared by:

ICF

Prepared for:

Center for Global Sustainability, University of Maryland

This section of the User Guide provides instruction on using the Land Use, Land-Use Change, and Forestry Module (LULUCF) module of the State Inventory Tool (SIT), and describes the methodology used for estimating greenhouse gas emissions (GHG) emissions and sinks from land use, land-use change, and forestry at the state level.

TABLE OF CONTENTS

1.1	Module Overview.....	3
1.1.1	Data Requirements	3
1.1.2	Tool Layout.....	4
1.2	Methodology.....	5
1.3	Uncertainty	23
1.4	References	23

1.1 MODULE OVERVIEW

This User Guide accompanies and explains the LULUCF module of the SIT.

Because most state inventories developed today rely heavily on the SIT, User Guides have been developed for each of the SIT modules. These modules contain largely updated methodologies that are, for the most part, consistent with the Inventory of U.S. Greenhouse Gas Emissions and Sinks (EPA 2025a) and related disaggregated data by state. Users can refer to the chapters and annexes of the U.S. Inventory to obtain additional information not found in the SIT or in the companion User Guide. Future updates will be consistent with the Center for Global Sustainability at University of Maryland's Greenhouse Gas Inventory and Analysis for the United States (GHGIA) (Desai et al. 2026).

In recent years, EPA also published an annual Inventory of U.S. Greenhouse Gas Emissions and Sinks by State (EPA 2024a) to make available consistent state-level GHG data for all states for use by states, researchers, and the general public. There will be differences between the latest EPA developed state-level estimates and inventory estimates developed by states using the SIT or other tools. Inventories compiled by states may differ for several reasons, and differences do not necessarily mean that one set of estimates is more accurate, or "correct". The CGS plans to continue EPA's initiative to publish an updated annual crosswalk guide to facilitate understanding methodological differences between methods in the current GHGIA, its disaggregation by state and SIT source categories, methods, and data.

The LULUCF module calculates CH₄ and N₂O emissions from fertilization of settlement soils, and forest fires, as well as carbon stock changes from forest management, urban trees, landfilled yard trimmings and food scraps, and agricultural soils.

The module provides default data for most inputs; however, if you have access to a more comprehensive or state-specific data source, it should be used in place of the default data. If using outside data sources, or for a more thorough understanding of the tool, please refer to the following discussion for data requirements and methodology.

1.1.1 Data Requirements


To calculate GHG emissions from land use, land-use change, and forestry, the data listed in Table 1 are required inputs (again, note that defaults are available for most of these data).

Table 1. Required data inputs for the LULUCF module.

Forestry Worksheets	Input Data Required
Forest Carbon Stock Change (Forest Land Remaining Forest Land, Land Converted to Forest Land, and Forest Land Converted to Land)	Carbon emitted from or sequestered in aboveground biomass, belowground biomass, dead wood, litter, mineral and organic soils, drained organic soils, wood products and landfills (million metric tons)
Urban Trees Carbon Stock Change	Carbon sequestration factor for urban trees (metric ton C/hectare/year) Total urban area (square kilometers) Urban area tree cover (percent)
N ₂ O from Settlement Soils	Direct N ₂ O emission factor for managed soils (percent) Total synthetic fertilizer applied to settlements (metric tons nitrogen)
Non-CO ₂ Emissions from Forest Fires	Emission factors for CH ₄ and N ₂ O emitted from burning forest and savanna (grams of gas/kilogram of dry matter combusted) Combustion efficiency of different vegetation types (percent) Average biomass density (kilograms dry matter per hectare) Area burned (hectares)
Landfilled Yard Trimmings and Food Scraps Carbon Stock Change	Grass, leaves, and branches constituting yard trimmings (percent) Yard trimmings and foods scraps landfilled, 1960-present (tons) Initial carbon content of yard trimmings and food scraps (percent) Dry weight/wet weight ratio of yard trimmings and foods scraps (percent) Proportion of carbon stored permanently for yard trimmings and foods scraps (percent) Half-life of degradable carbon for yard trimmings and foods scraps (years)
Agricultural Soil Carbon Stock Change	Carbon emitted from or sequestered in aboveground biomass, belowground biomass, dead wood, litter, and mineral and organic soils on cropland and grassland (million metric tons)

1.1.2 Tool Layout

Because there are multiple steps to complete within the LULUCF module, it is important to understand of the module's overall design. The layout of the LULUCF module and the purpose of its worksheets are presented in Figure 1.

Figure 1: Flow of Information in the LULUCF Module*

* These worksheets are the primary worksheets used in the LULUCF module; subsequent worksheets are used to populate the default data and are provided for informational purposes only.

1.2 METHODOLOGY

This section provides a guide to using the LULUCF module of the SIT to estimate GHG emissions and sequestration from land use, land-use change, and forestry. Within the LULUCF module, there are six sections: forest carbon flux (carbon stock change); urban trees; N₂O from settlement soils; non-CO₂ emissions from forest fires; carbon storage in landfilled yard trimmings and food scraps; and agricultural soil carbon flux. Because the methodology varies considerably among these sources/sinks, the details of each will be discussed in its respective step, following this general methodology discussion.

The LULUCF module follows the general methodology from the National Inventory Report (NIR) GHG Inventory by U.S. State. However, SIT aggregates changes in aboveground biomass, belowground biomass, deadwood, litter, and mineral and organic soil carbon stocks from Cropland Remaining Cropland, Land Converted to Cropland, Grassland Remaining Grassland, and Land Remaining Grassland under Ag Soil Carbon Flux in SIT. Default data is also not available for carbon flux from Wetlands Remaining Wetlands or from non-CO₂ emissions from Forest Fires; for these estimates, see the NIR GHG Inventory by U.S. State.

The LULUCF module will automatically calculate emissions after you enter the factors on the control worksheet and the required activity data on the individual sector worksheets. The tool provides default sector data for most sectors. The exception is forest fires where you will have to use an outside data source for area of forest burned per year (see Box 1 for suggested data sources).

Box 1: Forest Fire Data Sources

- Data are available from the National Interagency Coordination Center, which compiles fire records from Situation and Incident Status Summary (ICS-209) Reports. These records provide the number of acres burned by forest fire by state, and can be found in Table 12.4-2 of the EIP Guidance or online at https://www.nifc.gov/fireInfo/fireInfo_statistics.html
- To obtain the most accurate emission estimates, it is necessary to have information on the type of forests that have burned, as different types of forests contain differing amounts of combustible biomass. To further refine the analysis, information on specific burns and forest types can be found in the U.S. Federal Wildland Fire Management's website: <http://www.fs.fed.us/fire/>.
- To obtain accurate emissions for both wildfires and prescribed burning, users may directly consult FOFEM, which is available for download at <https://www.firelab.org/project/fofem>. Additional instructions for using the model are provided on the website.
- Land management agencies (e.g., the U.S. Forest Service, Bureau of Land Management, State Natural Resource Divisions) in each state maintain statistics on the areas and types of forests within their jurisdiction that have burned.
- State-level estimates of non-CO₂ emissions from forest fires are available for certain states from the U.S. Forest Service's publication, "[Greenhouse gas emissions and removals from forest land, woodlands, and urban trees in the United States, 1990-2023](#)" (see Appendix 1 of the publication). This information is also available in the 2025 NIR GHG Inventory by U.S. State released via Freedom of Information Act request (U.S. EPA 2025b).

There are nine general steps involved in estimating emissions using the LULUCF module: (1) select a state; (2) select an option for forest carbon flux; (3) enter emission factors and activity data for urban trees; (4) enter emission factors and activity data for N₂O from settlement soils; (5) enter emission factors and activity data for non-CO₂ emissions from forest fires; (6) enter emission factors and activity data for landfilled yard trimmings and food scraps; (7) select an option for agricultural soil carbon flux; (8) add any additional emissions sources that are not included in the module; and (9) review summary information. The general equations used to calculate GHG emissions from land use, land-use change, and forestry are provided below.

Step (1) Select a State

To begin, select the state you are interested in evaluating. By selecting a state, the rest of the tool will automatically reset to reflect the appropriate state default data and assumptions for use in subsequent steps of the tool.

Step (2) Select an Option for Forest Carbon Flux

Control Worksheet

The control worksheet allows you to either select the default data provided or to enter user-specified data to be used throughout the tool. To proceed with the default data, select the first radio button

under step 2 on the control worksheet. Default data are available for three land-use categories: *Forest Land Remaining Forest Land*, *Land Converted to Forest Land*, and *Forest Land Converted to Land*. These three categories are summed together on the Summary tab to calculate Net Forest Carbon Flux. Please note that a complete series of default data is not available for all states. If you would like to use your own state-specific data, select the second radio button under step 2 of the control worksheet. See Figure 2 for an example of the radio buttons in step 2.

Figure 2. Control Worksheet for the LULUCF Module

State Inventory Tool - Emissions and Sinks From Land Use, Land-Use Change, and Forestry

1. Choose a State: California (selected). **Choose a State** button. **Consult User's Guide** button. **Select All Defaults** button. **Reset All!** button. **Check/Uncheck All** button.

2. Forest Carbon Flux (Forest Land Remaining Forest Land, Land Converted to Forest Land, and Forest Land Converted to Land)

☐ Click here if you do not have any data on forest carbon flux and would like to use default estimates by the USDA Forest Service (no default data are available for AK, HI, or DC).

☒ Click here if you would like to use your own data on forest carbon flux. **Forest Carbon Flux Radio Buttons**

3.- 6. Enter emission factors and proceed to the corresponding worksheet to enter activity data for the following:

3. C Storage in Urban Trees

Carbon Sequestration Factor
metric ton C/hectare/year in California

Default Value: 2.88

4. N₂O from Settlement Soils

Emission Factor
Direct N₂O Emission Factor for Managed Soils

Default Value: 1%

5. Non-CO₂ From Forest Fires

Emission Factors

	Default Value	Value Used
Savanna, g CH ₄ /kg dry matter combusted	4.6	
Savanna, g N ₂ O/kg dry matter combusted	0.12	
Forests, g CH ₄ /kg dry matter combusted	8.1	
Forests, g N ₂ O/kg dry matter combusted	0.11	

Combustion Efficiency by Vegetation Type

	Default Value	Value Used
Primary tropical forests	36%	
Secondary tropical forests	55%	
Tertiary tropical forests	59%	
Boreal forest	34%	
Eucalypt forests	63%	
Other temperate forests	45%	

Individual Default Data Check Boxes

Go to Forest Carbon Flux Sheet

Go to Urban Trees Sheet

Go to Settlement Soils Sheet

Go to Burning CH₄ Sheet

Go to Burning N₂O Sheet

As a result of biological processes (e.g., growth and mortality) and anthropogenic activities (e.g., harvesting, thinning, and other removals), carbon is continuously cycled through ecosystem components, as well as between the forest ecosystem and the atmosphere. For example, the growth of trees results in the uptake of carbon from the atmosphere and storage in living trees. As these trees age, they continue to accumulate carbon until they reach maturity, at which point their carbon storage remains relatively constant. As trees die or drop branches and leaves on the forest floor, decay processes will release carbon to the atmosphere and also increase soil carbon. Some carbon from forests is also stored in wood products, such as lumber and furniture; and also in landfills, because when wood products are disposed of, they do not decay completely, and a portion of the carbon gets stored

indefinitely, as with landfilled yard trimmings and food scraps. The net change in forest carbon is the change in the amount of carbon stored in each of these pools (i.e., in each ecosystem component) over time. This section presents the methodology for calculating forest carbon flux/carbon stock change.

After completing the control worksheet for this sector, use the gray arrows to navigate to the *Forest Land Remaining Forest Land* worksheet.

Forest Land Remaining Forest Land Worksheet

If you are using the default data for forest carbon flux estimates, there is no further information to enter. Figure 3 shows the default worksheet for *Forest Land Remaining Forest Land*.

Figure 3. Example of Forest Land Remaining Forest Land Worksheet Using Default Data

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P	Q	R	S	T
	2a. Forest Land Remaining Forest Land in California																			
1	Two methodologies are used to calculate carbon emissions/storage (flux) from forest carbon using USDA Forest Service estimates of each state's forest carbon stocks.																			
2	(1) The first methodology applies to aboveground biomass, belowground biomass, dead wood, forest floor litter, and soil carbon from mineral and organic soils. USDA Forest Service estimates for each state's forest carbon stocks are provided for 1990-2018. These estimates are based on a publication titled "Greenhouse gas emissions and removals from forest land, woodlands, and urban trees in the United States, 1990-2018." Note: Net CO ₂ flux estimates presented in the table below are from the <i>Forest Land Remaining Forest Land</i> category. The following tabs provide estimates for <i>Land Converted to Forest Land</i> and <i>Forest Land Converted to Land</i> . These land-use change categories are combined on the Summary tab to provide an estimate of total forest carbon flux. No defaults are available for Alaska, Hawaii, or the District of Columbia. Totals may not sum to national or state-level totals due to independent rounding.																			
3	(2) The second methodology used applies to wood products and landfills (i.e. harvested wood products). Since the latest USDA Forest Service estimates for harvested wood products, default carbon emissions/storage are calculated by using USDA Forest Service estimates of each state's forest carbon stocks for 1990-2018. Changes from 1987-1992 and from 1992-1997 are each divided by 5 (the number of intervening years) to determine the average annual change for each year, giving total annual change. For the years 1998-2018, the average annual change for 1992-1997 is used as proxy data.																			
4	Users may also enter their own data. This may be done by selecting the appropriate option in Step 2 on the Control worksheet. For more information, see the Land Use Change, and Forestry chapter of the User's Guide.																			
5	Default data for Aboveground and Belowground Biomass, Dead Wood, Litter, and Soil Carbon																			
6	Net Sequestration/Emissions from Forest Land Remaining Forest Land																			
7	MMTCO ₂ E																			
8		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005			
9	Aboveground Biomass	(28.59)	(28.35)	(28.11)	(27.87)	(27.64)	(27.42)	(27.19)	(26.97)	(26.74)	(26.52)	(26.29)	(26.07)	(25.84)	(25.62)	(25.38)	(25.14)			
10	Belowground Biomass	(6.28)	(6.23)	(6.18)	(6.12)	(6.08)	(6.03)	(5.98)	(5.93)	(5.88)	(5.83)	(5.78)	(5.73)	(5.69)	(5.64)	(5.58)	(5.53)			
11	Deadwood	(5.68)	(5.72)	(5.76)	(5.79)	(5.82)	(5.84)	(5.87)	(5.89)	(5.91)	(5.94)	(5.96)	(5.99)	(6.01)	(6.04)	(6.04)	(6.04)			
12	Litter	(0.38)	(0.37)	(0.37)	(0.36)	(0.36)	(0.36)	(0.36)	(0.35)	(0.35)	(0.35)	(0.35)	(0.34)	(0.34)	(0.34)	(0.33)	(0.32)			
13	Soil (Mineral)	0.61	0.60	0.58	0.57	0.57	0.57	0.57	0.58	0.58	0.58	0.58	0.58	0.58	0.59	0.58	0.58			
14	Soil (Organic)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	-	-	-			
15	Drained Organic Soil	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-			
16	Total	(40.32)	(40.07)	(39.82)	(39.58)	(39.32)	(39.07)	(38.82)	(38.56)	(38.31)	(38.06)	(37.80)	(37.55)	(37.30)	(37.04)	(36.78)	(36.48)			

Next, click the gray navigational arrow to continue to the *Land Converted to Forest Land* worksheet.

Land Converted to Forest Land Worksheet

This tab calculates net sequestration/emissions from *Land Converted to Forest Land*. This category includes *Cropland Converted to Forest Land*, *Grassland Converted to Forest Land*, *Settlements Converted to Forest Land*, *Other Land Converted to Forest Land*, and *Wetlands Converted to Forest Land*. There is no further information to enter on this tab if you are using the default data for forest carbon flux. Figure 4 shows the default worksheet for *Land Converted to Forest Land*.

Figure 4: Example of Land Converted to Forest Land Worksheet Using Default Data

shown in Equation 1. The calculation is a sum of the fluxes for above- and belowground biomass, dead wood, litter, mineral and organic soils, drained organic soil, and wood products in use and in landfills. Once this sector worksheet is complete, use the gray navigational arrow to return to the control worksheet and proceed to the next step.

Equation 1. Forest Carbon Flux Equation

$$\text{Emissions or Sequestration (MMTCO}_2\text{E)} = \text{Sum of carbon fluxes from aboveground biomass, belowground biomass, dead wood, litter, mineral and organic soils, drained organic soil, and wood products and landfills}$$

Figure 6. Example of User-Entered Data Forest Carbon Flux Worksheet

2. Forest Carbon Flux in California

Click here to find possible data sources.

Enter net sequestration as a negative value, net emissions as a positive value in million metric tons of carbon dioxide equivalent. To use default data from USDA Forest Service, select the appropriate option in Step 2 on the control worksheet. Note: Default data are based on estimates of carbon flux from Forest Land Remaining, Forest Land, Land Converted to Forest Land, and Forest Land Converted to Land.

Return to Control Sheet

Clear All Data

Enter flux data in green cells

	Aboveground Biomass	Belowground Biomass	Dead Wood	Litter	Soil (Mineral)	Soil (Organic)	Total
	MMTCO ₂ E (million metric tons of carbon dioxide equivalent)						
1990							-
1991							-
1992							-
1993							-
1994							-
1995							-
1996							-
1997							-
1998							-
1999							-
2000							-
2001							-
2002							-
2003							-
2004							-
2005							-
2006							-
2007							-
2008							-
2009							-
2010							-
2011							-
2012							-

Step (3) Enter Emission Factors and Activity Data for Urban Trees

Control Worksheet

On the control worksheet, enter a carbon sequestration factor for urban trees in the appropriate yellow cell (metric tons of carbon per hectare per year). Default state-specific estimates of net sequestration are provided from Nowak et al. (2013). If the user-specific inputs do not match the default data in the control worksheet (i.e., the default value is overwritten), the text will appear red.

Trees in urban areas represent approximately 5.5 percent of total United States tree canopy cover (Nowak and Greenfield 2012). Furthermore, because trees in urban areas grow in relatively open surroundings, their growth and carbon sequestration are disproportionately large relative to forests. This section presents the methodology for calculating carbon sequestered by urban trees in your state.

After entering the appropriate sequestration factor, use the gray arrows to navigate to the Urban Trees worksheet.

Urban Trees Worksheet

Within the Urban Trees worksheet, enter data on the total urban area in your state (in square kilometers), as well as the average percent of urban area covered by trees, in the yellow cells. An example of this worksheet is shown in Figure 7. Equation 2 shows the method used to calculate carbon sequestration in urban trees.

Default urban areas are taken from Nowak et al. (2005) and the U.S. Census (1990, 2002, 2012, and 2023). Default state-specific percentages of urban tree cover are taken from Nowak and Greenfield (2012). Once this worksheet is complete, use the gray navigational arrow to return to the control worksheet and proceed to the next step.

Equation 2. Urban Trees Equation

$$\begin{aligned} \text{Sequestration (MMTCO}_2\text{E)} = & \\ & \text{Total Urban Area (km}^2\text{) x Urban Area with Tree Cover (\%)} \\ & \text{x 100 (ha/km}^2\text{) x C Sequestration Factor (metric tons C/ha/yr) x 44/12 (ratio of CO}_2\text{ to C) } \div \text{ 1,000,000 (to} \\ & \text{yield MMTCO}_2\text{E)} \end{aligned}$$

Figure 7. Example of Carbon Sequestration Factor Applied in the Urban Trees Worksheet

3. Urban Trees in California

Changes in carbon stocks in urban trees are equivalent to tree growth minus biomass losses resulting from pruning and mortality. Net carbon sequestration can be calculated using data on crown cover area or number of trees. Default state-specific data are given, or states may apply other state-specific values where available through sampling, aerial photography, or from municipal agencies that maintain urban vegetation data.

To estimate CO₂ sequestration by urban trees, the following steps are required: (1) obtain data on the area of urban tree cover; (2) calculate CO₂ flux; and (3) convert units to metric tons of carbon dioxide equivalent (MTCO₂E). This tool uses default urban area data multiplied by a state-specific estimate of the percent of urban area with tree cover to estimate the total area of urban tree cover. This default data, from Nowak et al. 2005, Nowak and Greenfield 2012, and the U.S. Census, represents urban area tree coverage for years 1990, 2000, and 2010. Estimates of urban area in the intervening years (1990-1999; 2001-2009) and subsequent years (2011-2016) are interpolated and extrapolated, respectively. State-specific net carbon sequestration rates are taken from Nowak et al. 2013 and multiplied by urban area to calculate CO₂ flux. States are encouraged to use state-specific data when available.

For more information, please consult the Land Use, Land-Use Change, and Forestry chapter of the User's Guide.

[Return to Control Sheet](#)

[Click here to find possible](#)

Required Data on Urban Area and Tree Cover

☒ Default Urban Area Data? [Clear All Data](#)

Year	Total Urban Area (km ²)	% of Urban Area with Tree Cover	Urban Tree Cover Area (km ²)	Carbon Sequestration Factor (metric ton C/hectare/year)	Carbon Sequestration (MMTCO ₂ E)
1990	17,600.00	25%	4,400.00	2.88	4.66
1991	17,898.40	25%	4,474.60	2.88	4.78
1992	18,196.80	25%	4,549.20	2.88	4.90
1993	18,495.20	25%	4,623.80	2.88	5.02
1994	18,793.60	25%	4,698.40	2.88	5.14
1995	19,092.00	25%	4,773.00	2.88	5.26
1996	19,390.40	25%	4,847.60	2.88	5.38
1997	19,688.80	25%	4,922.20	2.88	5.50
1998	19,987.20	25%	4,996.80	2.88	5.62
1999	20,285.60	25%	5,071.40	2.88	5.74

Carbon Sequestration Factor (from Control)

Step (4) Enter Emission Factors and Activity Data for N₂O from Settlement Soils

Control Worksheet

Of the fertilizers applied to soils in the United States, approximately 10 percent are applied to lawns, golf courses, and other landscaping occurring within settled areas. This section of the LULUCF module estimates N₂O emissions due to the application of fertilizers to settlement soils. On the control sheet you must enter an emission factor that will be used to calculate direct emissions due to fertilizer applications. The default value of 1 percent comes from IPCC (2006). If the user-specific inputs do not match the default data in the control worksheet (i.e., the default value is overwritten), the text will appear red.

After entering the appropriate emission factor, use the gray arrows to navigate to the Non-CO₂ from Settlement soils worksheet.

N₂O from Settlement Soils Worksheet

To complete this worksheet enter the amount of synthetic fertilizer applied (in metric tons of nitrogen) in the light blue cells (Figure 8). Emissions are calculated by multiplying the fertilizer data by the emission factor for direct emissions of N₂O (1.0 percent) to obtain the amount of emissions in N₂O-N/yr. This number is then converted from MT N₂O-N to MT N₂O by multiplying by the ratio of N₂O/N₂O-N (44/28). This is then converted to MMTCO₂E by dividing by 1,000,000 and multiplying by the GWP of N₂O. This calculation is shown in Equation 3. Once this worksheet is complete, use the gray navigational arrow to return to the control worksheet and proceed to emissions from forest fires.

Figure 8. Example of Fertilizer Data Applied in the Settlement Soils Worksheet

4. N₂O from Settlement Soils in California

Settlement soils include all developed land, including transportation infrastructure and human settlements of any size, unless they are already included under other categories.

N₂O emissions from settlement soils are calculated by multiplying the total synthetic fertilizer applied to settlements by the emission factor and the factor used to convert nitrogen to N₂O emissions (44/28). The calculated direct N₂O emissions are then multiplied by the global warming potential of N₂O and converted to million metric tons carbon dioxide equivalent.

For more information, please consult the User's Guide on estimating emissions from Land Use, Land-Use Change, and Forestry activities.

Click here to find possible data sources.

Return to Control Sheet

☒ Default Fertilizer Data?

	Required data on applied fertilizer (metric tons N)	Emission Factor (percent)	N ₂ O-N	Direct N ₂ O Emissions (metric tons N ₂ O-N/yr)	N ₂ O GWP	Carbon Dioxide Emissions (MTCO ₂ E)	Total Carbon Dioxide Emissions (MMTCO ₂ E)
1990	49,140	1%	1.57	723.6	298	230,114	0.230
1991	46,045	1%	1.57	723.6	298	215,624	0.216
1992	45,836	1%	1.57	720.3	298	214,642	0.215
1993	49,133	1%	1.57	772.1	298	230,085	0.230
1994	50,011	1%	1.57	785.9	298	234,194	0.234
1995	54,991	1%	1.57	864.1	298	257,513	0.258
1996	59,656	1%	1.57	937.5	298	279,360	0.279
1997	50,796	1%	1.57	798.2	298	237,871	0.238
1998	47,699	1%	1.57	749.5	298	223,366	0.223
1999	53,051	1%	1.57	833.7	298	248,430	0.248
2000	58,436	1%	1.57	918.3	298	273,647	0.274

Equation 3. Equation for Direct N₂O Emissions from Settlement Soils

$$\text{Emissions (MMTCO}_2\text{E)} = \text{Total Synthetic Fertilizer Applied to Settlement Soils (metric ton N)} \times \text{Emission Factor (percent)} \times 0.01 \text{ (metric tons N}_2\text{O-N/metric ton N)} \times 44/28 \text{ (Ratio of N}_2\text{O to N}_2\text{O-N)} \times 265 \text{ (GWP)} \div 1,000,000 \text{ (MT/MMTCO}_2\text{E)}$$

Step (5) Enter Emission Factors and Activity Data for Non-CO₂ from Forest Fires

Control Worksheet

On the control worksheet, the following data must be entered in the appropriate yellow cells for forest fires: (1) emission factors for N₂O and CH₄ for forest and savanna (grams of gas per kg dry matter combusted), (2) combustion efficiency by vegetation type (%), and (3) average biomass density in the state (kg dry matter per hectare). Default emission factors and combustion efficiencies are from IPCC (2006). Default biomass densities are adapted from Smith et al. (2001) and U.S. EPA (2003). States are encouraged to use this more detailed data if it is available and well documented. If the user-specific inputs do not match the default data in the control worksheet (i.e., the default value is overwritten), the text will appear red.

When a forest (or savanna) burns, the CO₂ emissions are included in the overall flux of forest carbon that is calculated in the forest carbon flux worksheet. However, forest fires also cause emissions of N₂O and CH₄ that are not accounted for under forest carbon flux because they are non-CO₂ emissions. This section presents the methodology for calculating N₂O and CH₄ emissions from forest fires.

After entering the appropriate emission factors for forest fires, use the gray arrows to navigate to the Non-CO₂ from Forest Fires worksheet.

Non-CO₂ from Forest Fires Worksheets

Within the Forest Fires worksheet, enter the area (hectares) burned per year in the pink cells. Because there is no default data available on area burned by state, you must rely on outside sources for this information (see Box 1 for suggestions). Equation 4 shows the method used to calculate N₂O and CH₄ emissions from forest fires. An example of this worksheet is shown in Figure 9. Once this sector worksheet is complete, use the gray navigational arrow to return to the control worksheet and proceed to the next category.

Equation 4. Forest Fires Emissions Equation

$$\text{Emissions (MMTCO}_2\text{E)} = \text{Area Burned (ha)} \times \text{Average Biomass Density (kg dry matter/ha)} \times \text{Combustion Efficiency (\%)} \times \text{Emission Factor (g gas/kg dry matter burned)} \times \text{GWP}$$

Figure 9. Example of Forest Fire Data Applied in the Forest Fire Worksheet

5a. Methane Emissions from Forest Fires in California

Biomass burned in forest fires emits CO₂, CH₄ and N₂O, in addition to many other gases and pollutants. CO₂ emissions are inherently captured under total carbon flux calculations, but CH₄ and N₂O must be estimated separately. All fires—wildfires and prescribed burns—emit these greenhouse gases.

Calculating the emissions of CH₄ and N₂O from burned forests requires determining the amount of carbon released by the fire (by multiplying the area burned, the fuel load, and the combustion efficiency) and then factoring in the emission ratio for each gas. No default data are available for area burned by forest type, but defaults are available for the other data points. States may choose to use state-specific data when available. The emission factors used in this tool are averages; to obtain more accurate data, users may wish to consult the National Fire Management Association's Guide.

Click here to find possible data sources.

Return to Control Sheet

Go to Burning N₂O Sheet

Clear All Data

Required data on area burned by forest fires

Combustion Efficiencies (from Control)

Emission Factors (from Control)

Forest Type	Area Burned (ha)	Average Biomass Density (kg d.m. / ha)	Combustion efficiency	Emission Factor (g/kg dry matter burned)	MTCH ₄ Emitted	CH ₄ GWP	Emissions MMTCO ₂ E
Primary tropical forests	x	150,061	x 36%	x 8.1	=	x 25	=
Secondary tropical forests	x	150,061	x 55%	x 8.1	=	x 25	=
Tertiary tropical forests	x	150,061	x 59%	x 8.1	=	x 25	=
Boreal forest	x	150,061	x 34%	x 8.1	=	x 25	=
Eucalypt forests	x	150,061	x 63%	x 8.1	=	x 25	=
Other temperate forests	x	150,061	x 45%	x 8.1	=	x 25	=
Shrublands	x	150,061	x 72%	x 8.1	=	x 25	=
Savanna woodlands (early dry season burns)	x	150,061	x 40%	x 4.6	=	x 25	=
Savanna woodlands (mid/late season burns)	x	150,061	x 74%	x 4.6	=	x 25	=
Total							

Biomass Density (from Control)

Forest Type	Area Burned (ha)	Average Biomass Density (kg d.m. / ha)	Combustion efficiency	Emission Factor (g/kg dry matter burned)	MTCH ₄ Emitted	CH ₄ GWP	Emissions MMTCO ₂ E
Primary tropical forests	x	150,061	x 36%	x 8.1	=	x 25	=
Secondary tropical forests	x	150,061	x 55%	x 8.1	=	x 25	=

Step (6) Enter Emission Factors and Activity Data for Landfilled Yard Trimming and Food Scraps

Control Worksheet

When wastes of biogenic origin (such as yard trimming and food scraps) are landfilled and do not completely decompose, the carbon that remains is stored. This section of the LULUCF module estimates the carbon stored in landfills by yard trimmings and food scraps.

Because the Landfilled Yard Trimmings and Food Scraps sector involves complicated calculations, the gray navigational arrow on the control worksheet takes you directly to the Landfilled Yard Trimmings and Food Scraps worksheet.

Landfilled Yard Trimmings and Food Scraps Worksheet

The Landfilled Yard Trimmings and Food Scraps sector worksheet calculates net carbon flux by estimating the change in landfilled carbon stocks between years, based on methodologies presented in IPCC (2003) and IPCC (2006). The LULUCF module uses Equation 5 to calculate carbon sequestration associated with landfilled yard trimmings and food scraps. Carbon stock estimates were calculated by: determining the mass of landfilled carbon resulting from yard trimmings or food scraps discarded in a given year; adding the accumulated landfilled carbon from previous years; and subtracting the portion of carbon landfilled in previous years that has decomposed.

Equation 5. Landfilled Yard Trimmings and Food Scraps Equation

$$LFC_{i,t} = \sum_n W_{i,n} \times (1 - MC_i) \times ICC_i \times \{ [CS_i \times ICC_i] + [(1 - (CS_i \times ICC_i)) \times e^{-k \times (t - n)}] \}$$

where,

- t** = the year for which carbon stocks are being estimated,
LFC_{i,t} = the stock of carbon in landfills in year t, for waste i (grass, leaves, branches, food scraps)
W_{i,n} = the mass of waste i disposed in landfills in year n, in units of wet weight
n = the year in which the waste was disposed, where 1960 < n < t
MC_i = moisture content of waste i,
CS_i = the proportion of initial carbon that is stored for waste i,
ICC_i = the initial carbon content of waste i,
e = the natural logarithm, and
k = the first order rate constant for waste i, and is equal to 0.693 divided by the half-life for decomposition.

To determine the total landfilled carbon stocks for a given year, the tool employs: (1) the composition of the yard trimmings; (2) the mass of yard trimmings and food scraps discarded in landfills; (3) the carbon storage factor of the landfilled yard trimmings and food scraps adjusted by mass balance; and (4) the rate of decomposition of the degradable carbon.

Due to the number of factors involved, the Landfilled Yard Trimmings and Food Scraps sector worksheet is arranged by a series of steps. To complete this sector worksheet, follow the steps below.

1. Enter the amount of landfilled yard trimmings and food scraps.
 - a. Enter the composition of yard trimmings in the orange cells as a percent of grass, leaves, and branches. Default percentages are available by clicking on the check boxes to the right of the orange input cells and are provided by Oshins and Block (2000). Figure 10 displays the inputs cells where this information is entered. If the user-specific inputs do not match the default data in the worksheet (i.e., the default value is overwritten), the text will appear red.
 - b. Enter the total annual landfilled yard trimmings and food scraps from 1960 to the present in short tons of wet weight in the yellow input cells. Default data are provided by clicking on the gray “Use default yard trimmings data” button above the yellow input cells. The tool uses the percentage entered for yard trimmings in the previous step to allocate the amount of yard trimmings distributed among grass, leaves, and branches. The default data from U.S. EPA (2020) is a national total for yard trimmings and food scraps and is distributed to each state based on state population U.S. Census Bureau (2026).

Figure 10. Landfilled Yard Trimmings and Food Scraps Worksheet, Step 1

6. Landfilled Yard Trimmings and Food Scraps in California

Estimates of net carbon flux of landfilled yard trimmings and food scraps can be calculated by estimating the change in landfill carbon stocks between inventory years. To determine the total landfilled carbon stocks for a given year, the following factors are estimated: (1) the composition of the yard trimmings, (2) the mass of yard trimmings and food scraps discarded in the state's landfills, (3) the carbon storage factor of the landfilled yard trimmings and food scraps, and (4) the rate of decomposition of the degradable carbon. The amount of carbon remaining in the landfill for each year is tracked based on a model of carbon fate that employs the equation outlined in Step 3 below.

Due to the complexity of these calculations, more detail about the methodology is provided below. Please note that many of the default factors are based on national values that may vary from state to state. States are encouraged to use state-specific data when available. For more information, please consult the Land Use, Land-Use Change, and Forestry Chapter of the User's Guide.

Click here to find possible data sources.

Return to Control Sheet

Clear All Data

1. Enter the composition of yard trimmings, and the amount of annually landfilled yard trimmings and food scraps.

Percent grass, leaves, and branches in yard trimmings

Content of yard trimmings	Default	Use the Default (Check for yes)
% Grass	30.3%	☐
% Leaves	40.1%	☐
% Branches	29.6%	☐

Check -- must add up to 100% in order to continue: 0% Must equal 100%

Landfilled yard trimmings and scraps, '000 short tons, wet weight

Default landfilled yard trimmings and food scraps = state population x national landfilled yard trimmings and food scraps

Default grass, leaves, and branches = total landfilled yard trimmings x percentages entered above

Use default yard trimmings data Clear Data

	1960	1961	1962	1963	1964	1969	1970	1971	1972
Total landfilled yard trimmings									
Grass									
Leaves									
Branches									
Food scraps									

Total landfilled yard trimmings and food scraps, 1960 to present

2. Calculate the amount of carbon added to landfills annually.

- Enter the initial carbon content percent for grass, leaves, branches, and food scraps in the orange cells, as shown in Figure 11. The default percentages are taken from Barlaz (1998). If the user-specific inputs do not match the default data in the worksheet (i.e., the default value is overwritten), the text will appear red.
- Enter the dry weight to wet weight ratio for grass, leaves, branches, and food scraps, also shown in Figure 13. This default information is drawn from Tchobanoglous, et al. (1993). If the user-specific inputs do not match the default data in the worksheet (i.e., the default value is overwritten), the text will appear red.

Figure 11. Landfilled Yard Trimmings and Food Scraps Worksheet, Step 2

6. Landfilled Yard Trimmings and Food Scraps in California

Estimates of net carbon flux of landfilled yard trimmings and food scraps can be calculated by estimating the change in landfill carbon stocks between inventory years. To determine the total landfilled carbon stocks for a given year, the following factors are estimated: (1) the composition of the yard trimmings, (2) the mass of yard trimmings and food scraps discarded in the state's landfills, (3) the carbon storage factor of the landfilled yard trimmings and food scraps, and 4) the rate of decomposition of the degradable carbon. The amount of carbon remaining in the landfill for each year is tracked based on a model of carbon fate that employs the equation outlined in Step 3 below.

Due to the complexity of these calculations, more detail about the methodology is provided below. Please note that many of the default factors are based on national values that may vary from state to state. States are encouraged to use state-specific data when available. For more information, please consult the Land Use, Land-Use Change, and Forestry Chapter of the User's Guide.

Click here to find possible data sources.

Return to Control Sheet

Clear All Data

2. Calculate the amount of carbon added to landfills annually

Key Assumptions

Initial Carbon Content	Default	Use the Default? (Check for Yes)
Grass	44.9%	☐
Leaves	45.5%	☐
Branches	49.4%	☐
Food Scraps	50.8%	☐

Enter initial carbon contents.

Enter dry weight to wet weight ratio

Dry Weight/Wet Weight ratio	Default	Use the Default? (Check for Yes)
Grass	30.0%	☐
Leaves	70.0%	☐
Branches	90.0%	☐
Food Scraps	30.0%	☐

Total Mass Additions, '000 metric tons C

Mass additions of carbon = landfilled materials, wet weight x initial carbon content x dry weight/wet weight ratio x metric tons per short ton

3. Calculate the total annual stocks of landfilled carbon.

- In the orange input cells, enter the proportion of carbon from each material stored in landfills indefinitely, as shown in Figure 12. Or use the default proportions, based on Barlaz (1998, 2005, and 2008). If the user-specific inputs do not match the default data in the worksheet (i.e., the default value is overwritten), the text will appear red.
- Enter the half-life of the degradable carbon in each of the materials in years, shown in Figure 14. The default data are from IPCC (2006). If the user-specific inputs do not match the default data in the worksheet (i.e., the default value is overwritten), the text will appear red.

Once this sector worksheet is complete, use the gray navigational arrow to return to the control worksheet.

Figure 12. Landfilled Yard Trimmings and Food Scraps Worksheet, Step 3

6. Landfilled Yard Trimmings and Food Scraps in California

Estimates of net carbon flux of landfilled yard trimmings and food scraps can be calculated by estimating the change in landfill carbon stocks between inventory years. To determine the total landfilled carbon stocks for a given year, the following factors are estimated: (1) the composition of the yard trimmings, (2) the mass of yard trimmings and food scraps discarded in the state's landfills, (3) the carbon storage factor of the landfilled yard trimmings and food scraps, and 4) the rate of decomposition of the degradable carbon. The amount of carbon remaining in the landfill for each year is tracked based on a model of carbon fate that employs the equation outlined in Step 3 below.

Due to the complexity of these calculations, more detail about the methodology is provided below. Please note that many of the default factors are based on national values that may vary from state to state. States are encouraged to use state-specific data when available. For more information, please consult the Land Use, Land-Use Change, and Forestry Chapter of the User's Guide.

[Click here to find possible data sources.](#)

[Return to Control Sheet](#)

[Clear All Data](#)

3. Calculate the total annual stocks of landfilled carbon

Proportion of Carbon Stored Permanently		Default	Use the Default (Check for Yes)	Use Default Percent for All?
Grass	53.5%	☐	☐	☐
Leaves	84.6%	☐	☐	☐
Branches	76.9%	☐	☐	☐
Food Scraps	15.7%	☐	☐	☐

Enter proportion of carbon stored permanently

Half-life of degradable carbon (years)		Default	Use the Default (Check for Yes)	Use Default Percent for All?
Grass	5	☐	☐	☐
Leaves	20	☐	☐	☐
Branches	23.1	☐	☐	☐
Food Scraps	3.8	☐	☐	☐

Enter half-life of degradable carbon

Total Stocks of Landfilled Carbon, '000 metric tons C

Annual carbon stocks are calculated by summing the carbon remaining from all previous years' deposits of waste.

The stock of carbon remaining in landfills from any given year is calculated as follows:

$$\text{Initial } C \text{ Addition} \times (\text{Proportion of } C \text{ Stored Permanently} + (1 - \text{Proportion of } C \text{ Stored Permanently}) \times e^{-\ln(0.5)/\text{half-life of degradable } C})$$

To calculate stocks for any given year, the remaining stocks for all previous years are summed.

The table below provides a summary of the calculated annual C stored in landfills. To view the more detailed calculations for each year, click on the navigational arrow below.

Step (7) Select an Option for Agricultural Soil Carbon Flux

Control Worksheet

The control worksheet allows you to either select the default data provided or to enter user-specified data to be used throughout the tool. To proceed with the default data, select the first radio button under step 7 on the control worksheet. If you would like to use your own state-specific data, select the second radio button under step 7 of the control worksheet. See Figure 13 for an example of the radio buttons in step 7.

Figure 13. Control Worksheet for the LULUCF Module

51

52 **7. Agricultural Soil Carbon Flux (in Cropland and Grassland soils)**

53 ☒ Click here if you do not have any data on carbon flux in agricultural soils and would like to use default estimates from the U.S. EPA's Inventory of Greenhouse Gas Emissions and Sinks: 1990-2015.

54 ☐ Click here if you would like to use your own data for agricultural soil carbon flux.

55

56 **8. Continue to estimate emissions . . .**

57

58 [Go to Summary Sheet](#)

59

60

61

62

63 **9. Export the results for use in the Synthesis Tool.**

64

65

66 [Export data](#)

67

68

Agricultural Soil Carbon Flux Radio Buttons

[Go to Agricultural Soil Carbon Flux Sheet](#)

Carbon is continuously cycled through the cropland and grassland ecosystems and the atmosphere. The amount of carbon stored in cropland varies according to crop type, management practices (e.g., rotation, tillage, drainage), and soil and climate variables. The amount of carbon stored in grassland depends on management practices (e.g., irrigation) and is also impacted by inter-annual climate variability, such as increased rainfall (IPCC 2006). Soil is the primary carbon pool in both cropland and grassland (U.S. EPA 2025a). Default data for agricultural soil carbon flux was updated in 2023 to include carbon flux from aboveground biomass, belowground biomass, deadwood, and litter in cropland and grassland ecosystems.

The net change in agricultural soil carbon is the change in the amount of carbon stored primarily in mineral and organic soils over time. This section presents the methodology for calculating agricultural soil carbon flux.

After completing the control worksheet for this sector, use the gray arrows to navigate to the Agricultural Soil Carbon Flux worksheet.

Agricultural Soil Carbon Flux Worksheet

If you are using the default data for agricultural soil carbon flux estimates, there is no further information to enter. Figure 14 shows the default agricultural soil carbon worksheet.

Figure 14. Example of Agricultural Soil Carbon Flux Worksheet Using Default Data

7. Agricultural Soil Carbon Flux in California

Default data for carbon emissions/storage (flux) from agricultural soils were obtained from U.S. EPA's *Inventory of Greenhouse Gas Emissions and Sinks: 1990-2018*. National estimates were apportioned to the state-level using a percentage breakdown reported from 2015. Agricultural soils include Cropland Remaining Cropland, Land Converted to Cropland, Grassland Remaining Grassland, and Land Converted to Grassland. Data for subsequent years (i.e., 2019-2020) were projected to 2018. Default data for Alaska and the District of Columbia are not available. Please refer to the *Inventory of Greenhouse Gas Emissions and Sinks: 1990-2018*, Chapter 6: Land Use, Land-Use Change, and Forestry for a summary of the methodology used to calculate carbon flux from agricultural soils.

Users may also enter their own data. This may be done by selecting the appropriate option in Step 7 on the Control worksheet. For more information, please consult the Land Use, Land-Use Change, and Forestry chapter of the User's Guide.

Return to Control Sheet

Default data for carbon stored in Cropland Remaining Cropland, Land Converted to Cropland, Grassland Remaining Grassland, and Land Converted to Grassland

Changes in Carbon Storage (million metric tons carbon)																
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999			2005	2006		
Total	0.945	1.076	0.917	0.578	1.268	0.818	1.376	1.015	1.194	1.402			1.947	2.246		

On default sheet, data are already provided

Net Sequestration/Emissions MMTCO ₂ E																	
	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006
Total	(3.463)	(3.945)	(3.364)	(2.118)	(4.649)	(3.001)	(5.046)	(3.723)	(4.377)	(5.139)	(7.511)	(7.042)	(7.900)	(7.289)	(7.002)	(7.140)	(8.234)

If you are using your own data on agricultural soil carbon flux, in the red cells enter total carbon flux data for cropland and grasslands (including land recently converted to cropland and grassland). Figure 15 shows the worksheet where you will enter agricultural soil carbon flux data. Once this sector worksheet is complete, use the gray navigational arrow to return to the control worksheet and proceed to the next step.

Figure 15. Example of User-Entered Data Agricultural Soil Carbon Flux Worksheet

7. Agricultural Soil Carbon Flux in California

Click here to find possible data sources.

Enter net sequestration as a negative value, net emissions as a positive value in million metric tons of carbon dioxide equivalent. To use default data from U.S. EPA's *Inventory of Greenhouse Gas Emissions and Sinks: 1990-2016*, select the appropriate option in Step 9 on the control worksheet.

Total		MMTCO ₂ E (million metric tons of carbon dioxide equivalent)	
1990	=		-
1991	=		-
1992	=		-
1993	=		-
1994	=		-
1995	=		-
1996	=		-
1997	=		-
1998	=		-
1999	=		-
2000	=		-
2001	=		-
2002	=		-
2003	=		-
2004	=		-
2005	=		-
2006	=		-

Enter flux data in red cells

Step (8) Add Additional Emissions Sources to Additional Sources Worksheet if Applicable

Users may use this worksheet to input data about additional emissions sources that are not included in this module or any other module. Additional emissions sources should be reported in MMTCO₂e. Figure 16 shows two options for entering data into this worksheet (please select only one):

Option A: Add up to ten additional emission sources in column B, lines 13-22. Enter the emissions in MMTCO₂e for each year 1990-2021 in the yellow cells.

Option B: Enter the CO₂, N₂O, and CH₄ (lines 33-35) for all other emissions sources for each year 1990-2021 in the yellow cells in MMTCO₂e.

Figure 16. Example Additional Emissions Sources Worksheet

8. Additional Emission Sources in Select a State . . .

Additional Emission Sources Data

Please use the "Additional Emission Sources" section of this sheet to enter any additional emission sources you would like to include in the inventory that are not captured in this or any other module. There are two options for entering data in this worksheet (please select only one):

Option A: Add up to ten additional emission sources in the column B, lines 13-22. Select the appropriate gas from the dropdown menu in column C. Enter the emissions in MMTCO₂e for each year 1990-2021 in the yellow cells.

Option B: Enter the total CO₂, CH₄, and N₂O (lines 33-35) for all other emissions sources for each year 1990-2021 in the yellow cells in MMTCO₂e.

For emission sources that do not have any default data or are not accounted for within the State Inventory Tool, state-level data are available here: <https://www.epa.gov/ghgemissions/state-ghg-emissions-and-removals>

[Return to Control Sheet](#)
[Clear all Data](#)

Additional Emission Sources (MMTCO₂E)

Option A

Source	Gas	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003

Option B

		1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
Additional Carbon Dioxide Emissions	CO ₂	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Additional Methane Emissions	CH ₄	-	-	-	-	-	-	-	-	-	-	-	-	-	-
Additional Nitrous Oxide Emissions	N ₂ O	-	-	-	-	-	-	-	-	-	-	-	-	-	-

For emission sources that do not have any default data or are not accounted for within the State Inventory Tool, state-level data are available here: <https://www.epa.gov/ghgemissions/state-ghg-emissions-and-removals>

Step (9) Review Summary Information

The steps above provide estimates of total emissions and sequestration from land use, land-use change, and forestry activities. The information from each sector worksheet is collected on the summary worksheet, which displays results in MMTCO₂E. Figure 17 shows the summary worksheet that sums the

emissions and sinks from all components of the LULUCF module. In addition, the results are displayed in graphical format at the bottom of the summary worksheets. Depending on whether the user entered their own data for forest carbon flux or used the default data provided by the module, additional rows may be shown to summarize emissions totals from *Forest Land Remaining Forest Land*, *Land Converted to Forest Land*, and *Forest Land Converted to Land*.

Figure 17. Example of the Emissions Summary Worksheet in the LULUCF Module

Return to Control Sheet

Review discussion of uncertainty associated with these results

Note: The Land Use, Land-Use Change, and Forestry module no longer estimates carbon dioxide emissions from Liming of Soils and Urea Fertilization. These categories are now estimated in the Agriculture module.

Emissions were not calculated for the following sector: Forest Fires. If you skipped any of these by mistake, please return to the control worksheet and complete each skipped source.

Emissions* (MMTCO ₂ E)	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001
Forest Carbon Flux	(54.54)	(54.29)	(54.05)	(50.04)	(49.78)	(49.53)	(49.27)	(49.02)	(48.77)	(48.51)	(48.26)	(48.01)
Aboveground Biomass	(28.59)	(28.35)	(28.11)	(27.87)	(27.64)	(27.42)	(27.19)	(26.97)	(26.74)	(26.52)	(26.29)	(26.07)
Belowground Biomass	(6.28)	(6.23)	(6.18)	(6.12)	(6.08)	(6.03)	(5.98)	(5.93)	(5.88)	(5.83)	(5.78)	(5.73)
Deadwood	(5.68)	(5.72)	(5.76)	(5.79)	(5.82)	(5.84)	(5.87)	(5.89)	(5.91)	(5.94)	(5.96)	(5.99)
Litter	(0.38)	(0.37)	(0.37)	(0.36)	(0.36)	(0.36)	(0.36)	(0.35)	(0.35)	(0.35)	(0.35)	(0.34)
Soil (Mineral)	0.61	0.60	0.58	0.57	0.57	0.57	0.57	0.58	0.58	0.58	0.58	0.58
Soil (Organic)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)	(0.00)
Drained Organic Soil	-	-	-	-	-	-	-	-	-	-	-	-
Total wood products and landfills	(14.22)	(14.22)	(14.22)	(10.46)	(10.46)	(10.46)	(10.46)	(10.46)	(10.46)	(10.46)	(10.46)	(10.46)
Urban Trees	(4.66)	(4.74)	(4.82)	(4.90)	(4.98)	(5.06)	(5.14)	(5.22)	(5.30)	(5.37)	(5.45)	(5.47)
Landfilled Yard Trimmings and Food Scraps	(2.94)	(2.80)	(2.77)	(2.42)	(2.12)	(1.74)	(1.44)	(1.53)	(1.52)	(1.43)	(1.42)	(1.46)
Grass	(0.24)	(0.22)	(0.22)	(0.17)	(0.14)	(0.10)	(0.06)	(0.06)	(0.07)	(0.06)	(0.06)	(0.07)
Leaves	(1.18)	(1.14)	(1.13)	(0.98)	(0.87)	(0.73)	(0.60)	(0.58)	(0.56)	(0.52)	(0.49)	(0.51)
Branches	(1.18)	(1.13)	(1.12)	(0.97)	(0.85)	(0.71)	(0.58)	(0.55)	(0.54)	(0.49)	(0.46)	(0.48)
Landfilled Food Scraps	(0.34)	(0.30)	(0.31)	(0.29)	(0.26)	(0.21)	(0.20)	(0.33)	(0.35)	(0.36)	(0.41)	(0.41)
Forest Fires	-	-	-	-	-	-	-	-	-	-	-	-
CH ₄	-	-	-	-	-	-	-	-	-	-	-	-
N ₂ O	-	-	-	-	-	-	-	-	-	-	-	-
N ₂ O from Settlement Soils	0.23	0.22	0.21	0.23	0.23	0.26	0.28	0.24	0.22	0.25	0.27	0.30
Agricultural Soil Carbon Flux	(3.46)	(3.95)	(3.36)	(2.12)	(4.65)	(3.00)	(5.05)	(3.72)	(4.38)	(5.14)	(7.51)	(7.04)
Total	(65.38)	(65.56)	(64.79)	(59.24)	(61.29)	(59.07)	(60.61)	(59.25)	(59.74)	(60.21)	(62.37)	(61.69)

* Note that parentheses indicate net sequestration.

1.3 UNCERTAINTY

In the upper right-hand corner of the summary worksheet is a button: “Review discussion of uncertainty associated with these results.” By clicking on this button, you are taken to a worksheet that discusses the uncertainty surrounding the activity data and emission factors, and how the uncertainty estimates for this source category affect the uncertainty of the emission estimates for your state.

1.4 REFERENCES

Barlaz, M.A. 2008. Corrections to Previously Published Carbon Storage Factors. In a letter report to Randy Freed of ICF International. February 28, 2008.

- Barlaz, M.A. 2005. "Decomposition of Leaves," letter report to Randy Freed of ICF International, June 29, 2005.
- Barlaz, M.A. 1998. "Carbon Storage during Biodegradation of Municipal Solid Waste Components in Laboratory-Scale Landfills." *Global Biogeochemical Cycles* 12: 373-380.
- Domke, Grant M.; Walters, Brian F.; Nowak, David J.; Smith, James, E.; Ogle, Stephen M.; Coulston, J.W.; Wirth, T.C. 2026. Greenhouse gas emissions and removals from forest land, woodlands, and urban trees in the United States, 1990-2023. Resource Update FS-382. Madison, WI: U.S. Department of Agriculture, Forest Service, Northern Research Station. 5 p.
<https://www.fs.usda.gov/rds/archive/catalog/RDS-2026-0031>.
- IPCC. 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Volume 4: Agriculture, Forestry, and Other Land Use. The National Greenhouse Gas Inventories Programme, The Intergovernmental Panel on Climate Change. [H.S. Eggleston, L. Buendia, K. Miwa, T. Ngara, and K. Tanabe (eds.)]. Hayama, Kanagawa, Japan.
- IPCC. 2003. *Good Practice Guidance for Land Use, Land-Use Change, and Forestry*. J. Penman and others, editors. IPCC National Greenhouse Gas Inventories Programme. Available online at <http://www.ipcc-nggip.iges.or.jp/public/gpglulucf/gpglulucf.htm>.
- M. Desai, V. Camobreco, T. Hedger, W. Irving, K. Rewcastle, J. Steller, L. Barbieri, M. Weitz, T. Murumkar, A. Fawcett, J. Lou, R. Cui, and N. Hultman. (2026). Greenhouse Gas Inventory and Analysis for the United States: 1990-2024. Center for Global Sustainability, University of Maryland, College Park.
- Nowak, D.J., E.J. Greenfield, R.E. Hoehn, and E. Lapoint. 2013. Carbon Storage and Sequestration by Trees in Urban and Community Areas of the United States. *Environmental Pollution* 178: 229-236. March 12, 2013.
- Nowak, D.J., E.J. Greenfield. 2018. Declining urban and community tree cover in the United States. *Urban Forestry & Urban Greening* (32) pp. 32-35. <https://www.fs.usda.gov/treesearch/pubs/55941>
- Nowak, D.J. and E.J. Greenfield. 2012. Tree and impervious cover in the United States. *Journal of Landscape and Urban Planning* (107) pp. 21-30.
- Nowak, D.J., J.T. Walton, L.G. Kaya, and J.F. Dwyer. 2005. "The Increasing Influence of Urban Environments on U.S. Forest Management." *Journal of Forestry* 103(8):377–382.
- Oshins, C., and D. Block. 2000. "Feedstock Composition at Composting Sites." *Biocycle* 41(9):31-34.
- Smith, J. 2014. Estimates of Forest Carbon Stocks and Flux: 1990-2014. E-mail correspondence between ICF and Jim Smith, USDA Forest Service. October 1, 2014 and January 27, 2015.
- Smith, W. Brad; Vissage, John S.; Darr, David R.; Sheffield, Raymond M. 2001. *Forest Resources of the United States, 1997*. Gen. Tech. Rep. NC-219. St. Paul, MN: United States Department of Agriculture, Forest Service, North Central Research Station. 190 p.

- Tchobanoglous, G., H. Theisen, and S.A. Vigil. 1993. *Integrated Solid Waste Management, 1st edition*. McGraw-Hill, NY. Cited by Barlaz (1998).
- U.S. Census Bureau. 2026. *State Population Totals and Components of Change: 2020-2025*. Available online at: <https://www.census.gov/data/tables/time-series/demo/popest/2020s-state-total.html>.
- U.S. Census Bureau. 2023. *A state-sorted list of all 2020 Census Urban Areas for the U.S., Puerto Rico, and Island Areas first sorted by state FIPS code, then sorted by Urban Area Census (UACE) code*. Available online at: <https://www.census.gov/programs-surveys/geography/guidance/geo-areas/urban-rural.html>
- U.S. Census Bureau. 2021. *National Population Totals: 2010-2020*. U.S. Census Bureau, Washington, DC. Available online at: <https://www.census.gov/programs-surveys/popest/technical-documentation/research/evaluation-estimates/2020-evaluation-estimates/2010s-totals-national.html>
- U.S. Census Bureau. 2012. Census 2010 Urbanized Area and Urban Cluster Information. Available at: https://www2.census.gov/geo/docs/reference/ua/ua_st_list_all.txt.
- U.S. Census Bureau. 2002. Census 2000 Urbanized Area and Urban Cluster Information. Available at: <https://www.census.gov/programs-surveys/geography/guidance/geo-areas/urban-rural/2000-urbanized-areas.html>.
- U.S. Census Bureau. 1990. Census 1990 Urbanized Area and Urban Cluster Information.
- U.S. EPA. 2003. *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990 – 2001*. Office of Atmospheric Protection, U.S. Environmental Protection Agency. Available at: <https://www.epa.gov/ghgemissions/inventory-us-greenhouse-gas-emissions-and-sinks-1990-2001>.
- U.S. EPA. 2020. *Advancing Sustainable Materials Management: Facts and Figures for 2018*. United States Environmental Protection Agency, Washington, DC. Available at <https://www.epa.gov/facts-and-figures-about-materials-waste-and-recycling/advancing-sustainable-materials-management>.
- U.S. EPA. 2024. Methodology report for inventory of U.S. greenhouse gas emissions and sinks by state: 1990–2022 (EPA-430-R-24-006). September. https://www.epa.gov/system/files/documents/2024-09/method-report_ghgibystate-508-september.pdf
- U.S. EPA. 2025a. Inventory of U.S. greenhouse gas emissions and sinks: 1990–2023 (EPA 430-R-25-003) [Unpublished annual report released via Freedom of Information Act Request]
- U.S. EPA. 2025b. DRAFT_AllStateGHGData_90-23. Unpublished Excel data file released via Freedom of Information Act request.